



UNITED STATES DEPARTMENT OF AGRICULTURE
WASHINGTON, D. C.

PRELIMINARY YIELD TABLES FOR SECOND-GROWTH STANDS IN THE CALIFORNIA PINE REGION

By DUNCAN DUNNING, *Silviculturist*, and L. H. REINEKE, *Associate Silviculturist*, California Forest and Range Experiment Station,¹ Branch of Research, Forest Service

CONTENTS

| | Page | | Page |
|---|------|--|------|
| Introduction..... | 1 | Yields—Continued. | |
| The forest types..... | 2 | Total basal area..... | 8 |
| Composition..... | 2 | Number of trees per acre..... | 11 |
| Range and occurrence..... | 2 | Average diameter of the stand..... | 13 |
| Conditions of establishment of second growth..... | 3 | Volume per acre in cubic feet..... | 14 |
| Relative importance of virgin and second-growth forest..... | 4 | Ratio of board feet to cubic feet..... | 16 |
| The data..... | 4 | Volume per acre in board feet..... | 18 |
| Methods used in analysis of data..... | 5 | Mean annual increment..... | 18 |
| Yields..... | 7 | Summary..... | 20 |
| Age..... | 7 | Literature cited..... | 21 |
| Site index (or quality)..... | 7 | Appendix..... | 21 |
| Density of stocking and stand-density index..... | 8 | Special technic used in the preparation of the tables..... | 21 |
| | | Statistical measures..... | 23 |

INTRODUCTION

The conifer forests of the California mountains are among the most valuable in North America. The character of the tree species forming the stands, the large dimensions attained by individual trees, the heavy yields, the extensive area occupied, and its comparative accessibility contribute to high values.

These high values have been hitherto associated almost entirely with virgin or old-growth stands. In recent years, however, second growth has begun to assume importance in the general scheme of forest economy. It is true that second growth has a present-utilization value in only rare instances of minor importance—the greater values still lie in the future; but for this very reason it has become desirable and is, indeed, becoming increasingly necessary to be able to forecast these values as accurately as possible.

Within the last 10 years two summaries of the forest situation, national in scope, have involved growth studies. In the management of the national forests a knowledge of second-growth yields is immediately useful in timber working-circle plans, in land exchange, in fire plans, in fire-damage appraisals, in cooperative fire-protection agreements with private landowners, and in other ways. To the

¹ Maintained in cooperation with the University of California.

State, yield data are of present value in formulating a land-use policy and in organizing fire protection for areas outside the national forests. Private timber owners may make use of yield tables in evaluating cut-over land for sale or exchange, in determining justifiable expenditures for fire protection, or in plans for continuous forest production. Such uses amply justify the attempt to put into convenient published form the data at present available, even though the complexity of yield-table construction for mixed stands and the difficulty in finding a sufficient number of adequate sample plots render these tables wholly tentative and preliminary in character.

THE FOREST TYPES

COMPOSITION

The stands to which the following tables apply are composed of ponderosa pine (*Pinus ponderosa* Laws.), sugar pine (*P. lambertiana* Dougl.), Douglas fir (*Pseudotsuga taxifolia* (Lam.) Britt.), white fir (*Abies concolor* Lindl. and Gord.), and red fir (*A. magnificia* A. Murray). Incense cedar (*Libocedrus decurrens* Torr.), although it also is almost invariably present (rare exceptions being in true fir stands) and frequently numerous, particularly in the younger age classes, rarely makes up more than 5 to 10 per cent of the volume of the stand.

The foregoing species appear in the stands in incalculably varying proportions, and yet, within clearly defined limits of altitude, latitude, and aspect, several types may be easily recognized. The types considered in this bulletin are shown in Table 1. The grouping of species into types is based on the arbitrary assumption that a species is a component of the type if it forms 20 per cent or more of the total basal area.

TABLE 1.—Composition of types, by species, in percentage of total basal area

| Species group or type | Ponderosa pine | Sugar pine | Douglas fir | White fir | Incense cedar | Red fir |
|------------------------------------|----------------|------------|-------------|-----------|---------------|----------|
| | Per cent | Per cent | Per cent | Per cent | Per cent | Per cent |
| Ponderosa pine-fir..... | 40 | 3 | 30 | 20 | 7 | |
| Ponderosa pine-sugar pine..... | 40 | 37 | 3 | 10 | 10 | |
| Ponderosa pine-sugar pine-fir..... | 40 | 25 | 10 | 20 | 5 | |
| Sugar pine-fir..... | 5 | 33 | 20 | 35 | 7 | |
| White fir-Douglas fir..... | 5 | 3 | 45 | 45 | 2 | |
| White fir-red fir..... | | 3 | | 68 | | 29 |

RANGE AND OCCURRENCE

The types here discussed cover the west slope of the Sierra Nevada and Cascade Mountains from the Kern River at the south to Mount Shasta at the north, and extend along the northern inner coast ranges from the Oregon line to central Lake County. The east-slope sierra and northeast plateau forests, where ponderosa and Jeffrey pines predominate, are not included in the discussion. The redwood and other forests of the northern outer coast ranges are also excluded.

The altitudinal limits of commercial stands vary with latitude and aspect. In the southern sierra good stands are rarely found below 3,000 feet, but in the northern inner coast ranges stands occur at elevations of 1,000 feet or less. The upper limit is about 6,500 feet in the south, and 5,000 feet in the north.

At lower elevations and on southerly aspects at middle elevations, ponderosa pine occurs in pure stands or predominates in mixtures. Toward the north, particularly in canyon bottoms and on north aspects, Douglas fir is frequently the dominant species at lower elevations. The sugar and ponderosa pine mixtures predominate on south slopes and benches between 4,000 and 5,500 feet. The sugar-pine and fir type is rather closely confined to north and northeast aspects at elevations of 4,500 to 6,000 feet. Above 6,000 feet, true firs usually predominate.

The best stands are found at elevations of 4,500 to 5,500 feet along the middle west slopes of the sierra. Here occurs the most favorable combination of precipitation and seasonal temperatures.

In relatively few sections of this large region are the stands uniform in age. All age classes, however, are rarely present, as they would be in a true selection forest. Stands are usually made up of small even-aged groups, the ages of the groups differing by periods of 10 to 20 years or more. The limited occurrence of extensive even-aged stands renders the construction and application of yield tables difficult.

CONDITIONS OF ESTABLISHMENT OF SECOND GROWTH

The occurrence of even-aged second-growth stands, within the limits fixed by nature, is largely a result of human activities. During the gold rush, beginning in 1849, heavy cuttings were made in the stands accessible to the mines. Frequently burning followed cutting, by accident or by design. The stands were rarely clear-cut at one time over considerable areas, but cutting was repeated, and a source of seed remained for some years. During seasons of heavy precipitation, such as the winter of 1861-62, the establishment of seedlings was good. The majority of the best even-aged second growth dates from the decade 1861-1870. Since most of the mining activity was at lower elevations, the greater portion of this second growth is ponderosa pine and Douglas fir.

During construction of the Southern Pacific Railroad through the Sierra, 1862 to 1865, heavy cutting along the right of way was followed by some excellent stands of second growth, varying in composition from pure ponderosa pine to pure fir.

Following the gold rush, stock raising increased rapidly, and range burning became a common practice. Although these range fires converted vast areas of forest into worthless brush, in many instances where conditions were favorable, patches of even-aged second growth were established.

With improved transportation the manufacture of lumber for the general market became important. The cuttings were light, only the more valuable pines being taken, as a rule. The resulting second growth is irregular in age.

Light cutting continued generally until the advent of heavier logging machinery and wider markets, beginning about 1900. From 1900 to about 1925, cutting on private land became more intensive, logging damage increased, and slash burning was more common. The frequency of fires, the failure of seed crops, and the generally poor climatic conditions have resulted in irregular, scanty reproduction for the most part. On national forest selective cuttings since 1906, the second growth is naturally irregular in age. Since 1925, private cutting has generally been lighter, leaving the firs, cedar, and smaller pines. Reproduction in these areas will not be even-aged.

Thus the conditions favoring the establishment of larger areas of even-aged second growth have largely ceased to exist. Probably the future tendency will be toward selection rather than even-aged stands. On the private land heavily cut or burned between 1900 and 1925, with better fire protection and more abundant precipitation, considerable even-aged reproduction may be expected.

RELATIVE IMPORTANCE OF VIRGIN AND SECOND-GROWTH FOREST

The areas of various classes of forest land in the region, as shown by the type map being prepared by the California Forest and Range Experiment Station, are given in Table 2. The area of virgin timber constitutes about 41 per cent of the area of all commercial timber in California, including redwood.

TABLE 2.—Areas of forested, restocking, and deforested lands in the California pine region

| Class of forest land | Area of class within pine region | |
|----------------------|----------------------------------|----------|
| | Acres | Per cent |
| Virgin timber..... | 4,663,400 | 47.8 |
| Second growth..... | 1,015,200 | 46.3 |
| Restocking..... | 869,100 | 49.7 |
| Deforested..... | 937,100 | 44.7 |

The virgin stands are not even-aged. The second growth includes all types of cut-over areas, and only a small percentage of it is even-aged. Restocking areas include mostly brush fields following early fires when the reproduction tends to be even-aged by groups and stocking is very irregular.

The future of the old-growth forests is, of course, uncertain. The pure ponderosa pine and the ponderosa pine-sugar pine types will doubtless be cut out first because of high value and accessibility. The comparatively small area of sugar pine-fir type will be culled for the sugar pine rather early. Types with a heavier proportion of Douglas fir and true firs will probably not be cut extensively until the available pine types are exhausted. It is to be expected that the privately owned stands will be largely cut over by the end of the century. They are more accessible, better stocked, and have a larger proportion of pines than the public forests.

The treatment of the remaining virgin timber will determine the future of second growth. So long as virgin timber is available, little cutting of second growth is to be expected. No cutting of consequence has occurred in young growth to date. Indications are that the ponderosa pine stands established 60 to 70 years ago will be the first second-growth utilized extensively.

THE DATA

The chief source of the material used in this study was a group of 224 plots measured by Dunning and Show (1912-1923). Permanent sample plots established later by the California Forest Experiment Station and other plots measured by F. X. Schumacher, then of the University of California, were also used. The total number of plots finally selected for the study was 311.

In several ways these plots were not entirely satisfactory. The outstanding feature is the irregular distribution by age. (Table 4.) The great preponderance of plots between the ages of 50 and 69 years is due, as has been shown, to the cuttings and frequent fires of the gold-mining period. Stands of other age classes are of relatively infrequent occurrence and hard to find.

Then, again, truly normal stands were so rare that, to obtain a sufficient number, a wider range of density of stocking than would ordinarily be permitted was unavoidable. Still further variation in stocking was doubtless due to the great variation in composition, which affected to some extent the judgment used in the selection of plots.

The field technic used when the earlier measurements were made by Dunning and Show differed in some details from the standard technic of the present. Curves of height over diameter were not prepared, and the height used for site classification was the average height of the dominant trees, rather than the height corresponding to the average diameter of the dominant and codominant trees. In addition, some of the plots were rather small, containing less than 100 trees, the present acceptable minimum. Despite these deficiencies, however, the data were of enough value to justify the construction of yield tables.

METHODS USED IN ANALYSIS OF DATA

The types themselves, in the relationship between component species, present a complex problem not encountered in the pure stands that are the basis of most of the recent work on second-growth yield tables. Although Haig's (5)² yield tables for western white pine in the northern Rocky Mountain region deal with a mixed-species forest, the associated species introduce no great complications since with increasing age they tend to disappear. No such tendency is noted in the California pine types, the component species of any young stand having equal likelihood of predominance at maturity.

This tendency to maintain the original composition throughout life requires, for the correct determination of site quality, an analysis of the interrelationship between species. It also requires an evaluation of the relative influence each species exerts upon such stand measurements as basal area, number of trees, etc. Such an evaluation, complicated by the endless variations in the proportions of the different species, ranging from almost pure stands of each species (except incense cedar) to any combination of two or more species, is complicated further by variations in density of stocking which tend to mask the effects of composition. Obviously, some method of measuring, defining, and compensating for stand density must be employed.

In general, the method of analysis consisted in assigning values for density of stocking to each plot and then modifying the stand measurements to represent the average of well-stocked stands. These modified values were used as a basis for constructing tables for average stocking and composition by the standard technic (3, 7). The percentage deviations of the original individual plot values from this average table were then correlated with density of stocking and

² Italic numbers in parentheses refer to Literature Cited, p. 21.

percentage composition by species. (Based on species distribution of basal area.) The resulting multiple regression equations gave the




FIGURE 1.—Site-classification curves, based upon the total height of the dominant tree of average basal area, of ponderosa pine, white fir, Douglas fir, and red fir, separately or in combination

appropriate factors to apply to the average table for any conditions of stocking and composition.

A detailed description of those particulars wherein the technic differed from standard procedure is given in the appendix.

YIELDS

AGE

The ages of the stands sampled were determined by ring counts on borings, usually at breast height, of dominant and codominant trees, corrected for the number of years required for the trees to reach the height at which the borings were made. The average age of the trees bored was taken as the age of the stand; no allowance was made for the period between the removal of the original stand and the establishment of the new stand.

SITE INDEX (OR QUALITY)

The classification of the sample plots according to site quality is based on the relationship between age and the average height of the dominant trees. Ponderosa pine, white fir, red fir, and Douglas fir have essentially the same height-age relationship when growing in mixed stands, and dominant trees of all the species named should be used as they are present.

Figure 1 and Table 3 show this relationship between age and height of dominant trees. The various site qualities are denoted by a site index expressed in height in feet attained at an age of 50 years. Obviously, a complete series of 1-foot site indices could be given but, for convenience, curves and tabular values are given only for even 10-foot intervals, from which intermediate values may be ascertained by interpolation. The determination of site index is simple; the average dominant height is plotted over the age on Figure 1, and the site index taken either from the curve nearest to the plotted point or from an interpolated index curve.

TABLE 3.—Total height in feet of average dominant tree ¹ on all sites

| Age (years) | Height on— | | | | | | | | | |
|-------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|----------------|----------------|
| | Site index 25 | Site index 30 | Site index 40 | Site index 50 | Site index 60 | Site index 70 | Site index 80 | Site index 90 | Site index 100 | Site index 110 |
| 20..... | 9.4 | 11.3 | 15.0 | 18.8 | 22.6 | 26.3 | 30.1 | 33.9 | 37.6 | 41.4 |
| 30..... | 14.6 | 17.5 | 23.4 | 29.2 | 35.1 | 40.9 | 46.8 | 52.6 | 58.4 | 64.3 |
| 40..... | 19.9 | 23.9 | 31.8 | 39.8 | 47.8 | 55.7 | 63.7 | 71.7 | 79.6 | 87.6 |
| 50..... | 25.0 | 30.0 | 40.0 | 50.0 | 60.0 | 70.0 | 80.0 | 90.0 | 100.0 | 110.0 |
| 60..... | 29.7 | 35.7 | 47.6 | 59.5 | 71.4 | 83.3 | 95.1 | 107 | 119 | 131 |
| 70..... | 34.0 | 40.8 | 54.3 | 67.9 | 81.5 | 95.1 | 109 | 122 | 136 | 149 |
| 80..... | 37.6 | 45.1 | 60.1 | 75.1 | 90.1 | 105 | 120 | 135 | 150 | 165 |
| 90..... | 40.6 | 48.7 | 64.9 | 81.1 | 97.4 | 114 | 130 | 146 | 162 | 178 |
| 100..... | 43.2 | 51.8 | 69.1 | 86.4 | 104 | 121 | 138 | 156 | 173 | 190 |
| 110..... | 45.6 | 54.7 | 72.9 | 91.1 | 109 | 128 | 146 | 164 | 182 | 201 |
| 120..... | 47.7 | 57.3 | 76.4 | 95.5 | 115 | 134 | 153 | 172 | 191 | 210 |
| 130..... | 49.8 | 59.7 | 79.7 | 99.6 | 119 | 139 | 159 | 179 | 199 | 219 |
| 140..... | 51.7 | 62.0 | 82.7 | 103 | 124 | 145 | 165 | 186 | 207 | 228 |
| 150..... | 53.5 | 64.2 | 85.6 | 107 | 128 | 150 | 171 | 193 | 214 | 236 |

¹ Based on dominant ponderosa pine, Douglas fir, red fir, and white fir; all four species to be used when present in the stand.

Table 4 shows how the 311 sample plots used are distributed as to age and site quality, and illustrates the preponderance of plots in the 50-59 year and 60-69 year age classes.

TABLE 4.—*Distribution of plots by age and site index*

| Age (years) | Number of plots on— | | | | | | | | | |
|-------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-------------------------|-----------|
| | Site indices 25-29 | Site indices 30-39 | Site indices 40-49 | Site indices 50-59 | Site indices 60-69 | Site indices 70-79 | Site indices 80-89 | Site indices 90-99 | Site indices 100-109 | All sites |
| 40 to 49 | | | | 2 | 7 | 5 | 4 | 1 | | 19 |
| 50 to 59 | 1 | 1 | 8 | 27 | 54 | 34 | 19 | 13 | 6 | 163 |
| 60 to 69 | | 1 | 10 | 25 | 26 | 17 | 4 | 4 | | 87 |
| 70 to 79 | | | 1 | | 3 | 3 | 3 | | | 10 |
| 80 to 89 | | | 1 | | 3 | | | | | 4 |
| 90 to 99 | | 1 | | 1 | 5 | 1 | 3 | | | 11 |
| 100 to 109 | | | | | 2 | | | | 1 | 3 |
| 110 to 119 | | | | 1 | 2 | 2 | 3 | | | 8 |
| 120 to 129 | | | | | 1 | 1 | | | | 2 |
| 130 to 139 | | | | | | 1 | | | | 1 |
| 140 to 149 | | | | 1 | | 2 | | | | 3 |
| All ages | 1 | 3 | 20 | 57 | 103 | 66 | 36 | 18 | 7 | 311 |

DENSITY OF STOCKING AND STAND-DENSITY INDEX

The criterion adopted for evaluating the density of the stand, or stocking, is the relationship between the number of trees per acre and their average diameter, shown by the solid line of Figure 2. This is a reference curve used for determining a stand-density index.³ The series of broken-line curves parallel to the reference curve serve a similar purpose to the curves in Figure 1 in supplying a ready means of estimating stand-density index. The number of trees per acre shown by each curve at an average diameter of 10 inches is the stand-density index of that curve. For a given stand, the line lying nearest the point defined by number of trees per acre plotted over average diameter can be read as the stand-density index. For example, the plotted point representing a stand with 350 trees per acre and an average diameter by basal area of 15 inches will fall nearest the curve representing a stand-density index of 700. By interpolating between the curves, a more precise index of 670 may be obtained.

The stand-density indices of the sample plots of this study ranged from 230 to 850, with an average value of 479. Taking 800 as a fair index for full stocking in these mixed-conifer stands, the average stand-density index of 479 represents approximately 60 per cent of full stocking. Individual plots ranged from 29 to 106 per cent. The distribution of the plots by stand-density index is given below:

| Stand-density index: | Number of plots |
|----------------------|-----------------|
| 200 to 299 | 6 |
| 300 to 399 | 49 |
| 400 to 499 | 102 |
| 500 to 599 | 80 |
| 600 to 699 | 50 |
| 700 to 799 | 18 |
| 800 to 899 | 6 |
| Total | 311 |

TOTAL BASAL AREA

Total basal area, or the sum of the breast-high cross-section areas of all trees 2 inches and larger in diameter is shown in Table 5 and Figure 3 for the average of well-stocked stands and average composi-

³ REINEKE, L. H. PERFECTING A STAND-DENSITY INDEX FOR EVEN-AGED STANDS. (Unpublished manuscript.)




FIGURE 2.—Stand-density index curves. When, for a given stand, trees per acre over average diameter by basal area is plotted, the stand-density index is determined in round numbers by the nearest curve, or more exactly by interpolation

158797°-33-2

tion. The modification of this table for any condition of stocking and composition is effected by substituting the appropriate values in the regression equation given under the table and applying the resulting percentage to the values given in Table 5.




FIGURE 3.—Relationship between age, site index, and total basal area for average well-stocked stands of average composition

For convenience, the percentage corrections for various stand-density indices, by types, are given in Table 6. Minor differences in composition from that given for these types will not change the correction percentage appreciably.

TABLE 5.—Total basal area, average stands ¹

| Age (years) | Basal area (square feet per acre) by site-index classes | | | | | | | | | |
|-------------|---|---------|---------|---------|---------|---------|---------|---------|----------|----------|
| | Site 25 | Site 30 | Site 40 | Site 50 | Site 60 | Site 70 | Site 80 | Site 90 | Site 100 | Site 110 |
| 30..... | 124 | 126 | 131 | 138 | 146 | 155 | 164 | 172 | 178 | 184 |
| 40..... | 165 | 168 | 175 | 184 | 196 | 208 | 219 | 229 | 238 | 245 |
| 50..... | 194 | 197 | 205 | 216 | 230 | 244 | 257 | 269 | 280 | 288 |
| 60..... | 210 | 213 | 222 | 234 | 248 | 264 | 278 | 291 | 303 | 311 |
| 70..... | 223 | 226 | 236 | 248 | 264 | 280 | 295 | 309 | 321 | 331 |
| 80..... | 234 | 237 | 247 | 260 | 277 | 294 | 310 | 324 | 337 | 347 |
| 90..... | 243 | 247 | 257 | 271 | 288 | 305 | 322 | 337 | 350 | 361 |
| 100..... | 252 | 255 | 266 | 280 | 298 | 316 | 333 | 349 | 363 | 373 |
| 110..... | 258 | 262 | 273 | 287 | 305 | 324 | 342 | 358 | 372 | 383 |
| 120..... | 264 | 268 | 279 | 294 | 312 | 332 | 350 | 366 | 381 | 392 |
| 130..... | 269 | 273 | 285 | 300 | 319 | 338 | 357 | 374 | 388 | 400 |
| 140..... | 274 | 278 | 290 | 305 | 324 | 344 | 363 | 380 | 395 | 407 |
| 150..... | 278 | 282 | 294 | 310 | 329 | 350 | 369 | 386 | 401 | 413 |

¹ For specific stands, substitute in the following equation and apply resulting percentage to the tabular values: Total basal area (in percentage of composite table) = 2.0303 (percentage stocking) + 0.1493 (ponderosa pine per cent) + 0.1545 (sugar pine per cent) + 0.0541 (Douglas fir per cent) + 0.0980 (white fir per cent) + 0.1229 (incense cedar per cent) + 0.1883 (red fir per cent) - 8.1511.

TABLE 6.—Percentage correction of values in Table 5 for different types ¹

| Stand-density index | Ponderosa pine-fir | Ponderosa pine-sugar pine | Ponderosa pine-sugar pine-fir | Sugar pine-fir | White fir-Douglas fir | White fir-red fir |
|---------------------|--------------------|---------------------------|-------------------------------|----------------|-----------------------|-------------------|
| | Per cent | Per cent | Per cent | Per cent | Per cent | Per cent |
| 200..... | 43.33 | 46.51 | 45.40 | 43.67 | 40.76 | 37.92 |
| 250..... | 53.49 | 56.67 | 55.56 | 53.32 | 50.91 | 48.07 |
| 300..... | 63.64 | 66.82 | 65.71 | 63.98 | 61.06 | 58.22 |
| 350..... | 73.79 | 76.97 | 75.86 | 74.13 | 71.21 | 68.37 |
| 400..... | 83.94 | 87.12 | 86.01 | 84.28 | 81.36 | 78.52 |
| 450..... | 94.09 | 97.27 | 96.16 | 94.43 | 91.51 | 88.67 |
| 500..... | 104.24 | 107.42 | 106.31 | 104.58 | 101.66 | 98.82 |
| 550..... | 114.39 | 117.58 | 116.47 | 114.73 | 111.82 | 108.98 |
| 600..... | 124.55 | 127.73 | 126.62 | 124.88 | 121.97 | 119.13 |
| 650..... | 134.70 | 137.88 | 136.77 | 135.04 | 132.12 | 129.28 |
| 700..... | 144.85 | 148.03 | 146.92 | 145.19 | 142.27 | 139.43 |
| 750..... | 155.00 | 158.18 | 157.07 | 155.34 | 152.42 | 149.58 |
| 800..... | 165.15 | 168.33 | 167.22 | 165.49 | 162.57 | 159.73 |

¹ Aggregate deviation, less than 0.5 per cent; average percentage deviation, 5.22 per cent.

NUMBER OF TREES PER ACRE

The number per acre of all trees 2 inches or more in breast-high diameter is given in Table 7 and Figure 4 for average stocking and composition. Modification for specific stocking and composition is made as for total basal area. Table 8 gives the percentages for the six types.




FIGURE 4.—Relationship between age, site index, and number of trees per acre for average well-stocked stands of average composition

TABLE 7.—Number of trees, average stands ¹

| Age (years) | Number of trees per acre, by site index class | | | | | | | | | |
|-------------|---|---------|---------|---------|---------|---------|---------|---------|----------|----------|
| | Site 25 | Site 30 | Site 40 | Site 50 | Site 60 | Site 70 | Site 80 | Site 90 | Site 100 | Site 110 |
| 30..... | 3,260 | 3,030 | 2,620 | 2,175 | 1,770 | 1,400 | 1,120 | 922 | 790 | 690 |
| 40..... | 2,110 | 1,960 | 1,695 | 1,405 | 1,145 | 906 | 723 | 596 | 511 | 446 |
| 50..... | 1,500 | 1,395 | 1,205 | 1,000 | 814 | 645 | 515 | 424 | 364 | 318 |
| 60..... | 1,095 | 1,020 | 879 | 730 | 594 | 470 | 375 | 309 | 265 | 232 |
| 70..... | 807 | 750 | 648 | 538 | 438 | 347 | 277 | 228 | 196 | 171 |
| 80..... | 648 | 502 | 520 | 432 | 351 | 278 | 222 | 183 | 157 | 137 |
| 90..... | 547 | 509 | 439 | 365 | 297 | 235 | 188 | 155 | 133 | 116 |
| 100..... | 481 | 447 | 386 | 321 | 261 | 207 | 165 | 136 | 117 | 102 |
| 110..... | 437 | 406 | 351 | 291 | 237 | 188 | 150 | 123 | 106 | 92 |
| 120..... | 403 | 374 | 323 | 268 | 218 | 173 | 138 | 114 | 98 | 85 |
| 130..... | 377 | 350 | 303 | 251 | 204 | 162 | 129 | 107 | 91 | 80 |
| 140..... | 356 | 331 | 286 | 237 | 193 | 153 | 122 | 101 | 86 | 75 |
| 150..... | 337 | 313 | 270 | 224 | 182 | 145 | 115 | 95 | 82 | 71 |

¹ For specific stands, substitute in the following equation and apply resulting percentage to the tabular values: Number of trees (in percentage of composite table) = 2.0058 (percentage of stocking) - 0.3166 (ponderosa pine per cent) - 0.2973 (sugar pine per cent) - 0.0159 (Douglas fir per cent) - 0.1355 (white fir per cent) - 0.1358 (incense cedar per cent) - 0.0905 (red fir per cent) + 19.3572.

TABLE 8.—Percentage correction of values in Table 7 for different types ¹

| Stand-density index | Ponderosa pine-fir | Ponderosa pine-sugar pine | Ponderosa pine-sugar pine-fir | Sugar pine-fir | White fir-Douglas fir | White fir-red fir |
|---------------------|--------------------|---------------------------|-------------------------------|----------------|-----------------------|-------------------|
| | Per cent | Per cent | Per cent | Per cent | Per cent | Per cent |
| 200..... | 41.78 | 33.05 | 35.83 | 42.07 | 49.91 | 46.74 |
| 250..... | 51.81 | 43.08 | 45.86 | 52.10 | 59.94 | 56.77 |
| 300..... | 61.84 | 53.11 | 55.89 | 62.13 | 69.97 | 66.80 |
| 350..... | 71.87 | 63.14 | 65.92 | 72.16 | 80.00 | 76.83 |
| 400..... | 81.90 | 73.16 | 75.94 | 82.18 | 90.03 | 86.86 |
| 450..... | 91.92 | 83.19 | 85.97 | 92.21 | 100.06 | 96.89 |
| 500..... | 101.95 | 93.22 | 96.00 | 102.24 | 110.09 | 106.92 |
| 550..... | 111.98 | 103.25 | 106.03 | 112.27 | 120.12 | 116.95 |
| 600..... | 122.01 | 113.28 | 116.06 | 122.30 | 130.15 | 126.97 |
| 650..... | 132.04 | 123.31 | 126.09 | 132.33 | 140.17 | 137.00 |
| 700..... | 142.07 | 133.34 | 136.12 | 142.36 | 150.20 | 147.03 |
| 750..... | 152.10 | 143.37 | 146.15 | 152.39 | 160.23 | 157.06 |
| 800..... | 162.13 | 153.40 | 156.18 | 162.42 | 170.26 | 167.09 |

¹ Aggregate deviation, less than 0.5 per cent; average percentage deviation, 18.4 per cent.

AVERAGE DIAMETER OF THE STAND

The average breast-high diameter of the stand is the diameter of the tree of average basal area as determined by dividing total basal area by number of trees. Table 9 and Figure 5 give these values for stands of average stocking and composition. For stands of other than average stocking or composition, average breast-high diameter should be derived by dividing the corrected total basal area by the corrected number of trees and converting the resultant average basal area into its equivalent diameter.

TABLE 9.—Average breast-high diameter in inches of average stands, by site-index classes

| Age (years) | Site 25 | Site 30 | Site 40 | Site 50 | Site 60 | Site 70 | Site 80 | Site 90 | Site 100 | Site 110 |
|-------------|---------|---------|---------|---------|---------|---------|---------|---------|----------|----------|
| 30 | 2.64 | 2.76 | 3.03 | 3.41 | 3.89 | 4.51 | 5.18 | 5.85 | 6.43 | 6.99 |
| 40 | 3.79 | 3.96 | 4.35 | 4.90 | 5.60 | 6.49 | 7.45 | 8.39 | 9.24 | 10.0 |
| 50 | 4.87 | 5.09 | 5.59 | 6.29 | 7.20 | 8.33 | 9.56 | 10.8 | 11.9 | 12.9 |
| 60 | 5.93 | 6.19 | 6.81 | 7.67 | 8.75 | 10.1 | 11.7 | 13.1 | 14.5 | 15.7 |
| 70 | 7.12 | 7.43 | 8.17 | 9.19 | 10.5 | 12.2 | 14.0 | 15.8 | 17.3 | 18.8 |
| 80 | 8.14 | 8.50 | 9.33 | 10.5 | 12.0 | 13.9 | 16.0 | 18.0 | 19.8 | 21.5 |
| 90 | 9.02 | 9.43 | 10.4 | 11.7 | 13.3 | 15.4 | 17.7 | 20.0 | 22.0 | 23.9 |
| 100 | 9.80 | 10.2 | 11.2 | 12.6 | 14.5 | 16.7 | 19.2 | 21.7 | 23.9 | 25.9 |
| 110 | 10.4 | 10.9 | 11.9 | 13.5 | 15.4 | 17.8 | 20.4 | 23.1 | 25.4 | 27.6 |
| 120 | 11.0 | 11.5 | 12.6 | 14.2 | 16.2 | 18.8 | 21.6 | 24.3 | 26.7 | 29.0 |
| 130 | 11.4 | 12.0 | 13.1 | 14.8 | 16.9 | 19.6 | 22.5 | 25.3 | 28.0 | 30.3 |
| 140 | 11.9 | 12.4 | 13.6 | 15.4 | 17.5 | 20.3 | 23.4 | 26.3 | 29.0 | 31.5 |
| 150 | 12.3 | 12.9 | 14.2 | 15.9 | 18.2 | 21.0 | 24.3 | 27.3 | 29.9 | 32.7 |




FIGURE 5.—Relationship between age, site index, and average diameter per acre, for average well-stocked stands of average composition

VOLUME PER ACRE IN CUBIC FEET

Table 10 and Figure 6 show for stands of average stocking and composition the total cubic-foot volume inside bark, including stump and top, of all trees 2 inches and larger in diameter. Correction percentages are given in Table 11 for the six types.




FIGURE 6.—The volumes in cubic feet (inside bark, stump and top included, for average well-stocked stands of average composition

TABLE 10.—Volume per acre in cubic feet, average stands ¹

| Age (years) | Volume in cubic feet per acre on site index— | | | | | | | | | |
|-------------|--|-------|--------|--------|--------|--------|--------|--------|--------|--------|
| | 25 | 30 | 40 | 50 | 60 | 70 | 80 | 90 | 100 | 110 |
| 30..... | 930 | 1,060 | 1,420 | 1,800 | 2,200 | 2,650 | 3,200 | 3,770 | 4,380 | 4,850 |
| 40..... | 1,590 | 1,800 | 2,410 | 3,090 | 3,770 | 4,550 | 5,500 | 6,500 | 7,450 | 8,450 |
| 50..... | 2,260 | 2,590 | 3,480 | 4,440 | 5,420 | 6,560 | 7,940 | 9,300 | 10,700 | 12,000 |
| 60..... | 2,950 | 3,350 | 4,510 | 5,790 | 7,020 | 8,550 | 10,300 | 12,100 | 13,900 | 15,500 |
| 70..... | 3,500 | 4,000 | 5,350 | 6,850 | 8,450 | 10,200 | 12,200 | 14,300 | 16,500 | 18,400 |
| 80..... | 4,000 | 4,540 | 6,100 | 7,800 | 9,520 | 11,500 | 13,800 | 16,200 | 18,600 | 20,900 |
| 90..... | 4,420 | 5,000 | 6,750 | 8,600 | 10,500 | 12,700 | 15,200 | 17,800 | 20,700 | 23,100 |
| 100..... | 4,830 | 5,500 | 7,400 | 9,450 | 11,500 | 13,900 | 16,600 | 19,500 | 22,600 | 25,200 |
| 110..... | 5,200 | 5,950 | 8,000 | 10,200 | 12,300 | 14,900 | 17,900 | 21,000 | 24,250 | 27,200 |
| 120..... | 5,600 | 6,390 | 8,550 | 10,800 | 13,200 | 16,000 | 19,100 | 22,500 | 26,000 | 29,000 |
| 130..... | 5,900 | 6,700 | 9,050 | 11,500 | 14,000 | 16,900 | 20,300 | 24,000 | 27,500 | 30,700 |
| 140..... | 6,220 | 7,070 | 9,550 | 12,100 | 14,750 | 17,750 | 21,400 | 25,200 | 29,000 | 32,400 |
| 150..... | 6,550 | 7,430 | 10,000 | 12,700 | 15,450 | 18,550 | 22,400 | 26,400 | 30,400 | 34,000 |

¹ For specific stands, substitute in the following equation and apply resulting percentage to the tabular values: Volume in cubic feet (in percentage of composite table) = 1.5251 (percentage of stocking) - 0.7170 (ponderosa pine per cent) - 0.7740 (sugar pine per cent) - 0.9398 (Douglas fir per cent) - 1.0704 (white fir per cent) - 0.8427 (incense cedar per cent) - 1.1936 (red fir per cent) + 123.7553.

TABLE 11.—Percentage correction of Table 10 for different types ¹

| Stand density index | Ponderosa pine-fir | Ponderosa pine-sugar pine | Ponderosa pine-sugar pine-fir | Sugar pine-fir | White fir-Douglas fir | White fir-red fir |
|---------------------|--------------------|---------------------------|-------------------------------|-----------------|-----------------------|-------------------|
| | <i>Per cent</i> | <i>Per cent</i> | <i>Per cent</i> | <i>Per cent</i> | <i>Per cent</i> | <i>Per cent</i> |
| 200..... | 74.01 | 79.99 | 76.21 | 67.97 | 61.21 | 49.36 |
| 250..... | 81.64 | 87.61 | 83.83 | 75.60 | 68.83 | 56.99 |
| 300..... | 89.27 | 95.24 | 91.46 | 83.22 | 76.46 | 64.61 |
| 350..... | 96.89 | 102.87 | 99.08 | 90.85 | 84.08 | 72.24 |
| 400..... | 104.52 | 110.49 | 106.71 | 98.47 | 91.71 | 79.86 |
| 450..... | 112.14 | 118.12 | 114.34 | 106.10 | 99.33 | 87.49 |
| 500..... | 119.77 | 125.74 | 121.96 | 113.72 | 106.96 | 95.11 |
| 550..... | 127.39 | 133.37 | 129.59 | 121.35 | 114.58 | 102.74 |
| 600..... | 135.02 | 140.99 | 137.21 | 128.98 | 122.21 | 110.36 |
| 650..... | 142.64 | 148.62 | 144.84 | 136.60 | 129.84 | 117.99 |
| 700..... | 150.27 | 156.24 | 152.46 | 144.23 | 137.46 | 125.61 |
| 750..... | 157.89 | 163.87 | 160.09 | 151.85 | 145.09 | 133.24 |
| 800..... | 165.52 | 171.49 | 167.71 | 159.48 | 152.71 | 140.87 |

¹ Aggregate deviation, less than 0.5 per cent; average percentage deviation, 16.4 per cent.




FIGURE 7.—The ratio of volume of stand in board feet to volume in cubic feet, in relation to average diameter of stand

RATIO OF BOARD FEET TO CUBIC FEET

The ratio of board feet to cubic feet is closely correlated with average diameter and is independent of age and site quality. In the standard yield-table technic, this relationship is applied to the table

of volume in cubic feet to obtain the table of volume in board feet. Table 12 and Figure 7 give the board foot-cubic foot ratios used. The values given are ratios of volume in board feet of trees 8 inches and larger to the volume in cubic feet of all trees 2 inches and larger.




FIGURE 8.—Volume in board feet by the international rule ($\frac{1}{8}$ -inch kerf), trees 8 inches and larger, average of well-stocked stands, average composition.

TABLE 12.—Board-foot conversion table

| Age (years) | Board feet per cubic foot by site-index classes | | | | | | | | | |
|-------------|---|---------|---------|---------|---------|---------|---------|---------|----------|----------|
| | Site 25 | Site 30 | Site 40 | Site 50 | Site 60 | Site 70 | Site 80 | Site 90 | Site 100 | Site 110 |
| 30 | | | 0.06 | 0.25 | 0.50 | 0.80 | 1.20 | 1.60 | 1.98 | 2.29 |
| 40 | 0.44 | 0.52 | .72 | 1.04 | 1.47 | 2.00 | 2.60 | 3.15 | 3.60 | 3.96 |
| 50 | 1.02 | 1.15 | 1.45 | 1.87 | 2.44 | 3.10 | 3.76 | 4.31 | 4.72 | 5.06 |
| 60 | 1.60 | 1.82 | 2.21 | 2.72 | 3.35 | 4.01 | 4.65 | 5.12 | 5.52 | 5.82 |
| 70 | 2.38 | 2.56 | 2.98 | 3.55 | 4.19 | 4.80 | 5.36 | 5.82 | 6.19 | 6.46 |
| 80 | 2.98 | 3.20 | 3.60 | 4.19 | 4.76 | 5.36 | 5.90 | 6.32 | 6.60 | 6.81 |
| 90 | 3.50 | 3.70 | 4.14 | 4.65 | 5.17 | 5.75 | 6.30 | 6.65 | 6.87 | 7.01 |
| 100 | 3.85 | 4.05 | 4.47 | 4.96 | 5.49 | 6.06 | 6.52 | 6.84 | 7.00 | 7.12 |
| 110 | 4.14 | 4.35 | 4.76 | 5.24 | 5.75 | 6.30 | 6.71 | 6.96 | 7.10 | 7.19 |
| 120 | 4.39 | 4.58 | 4.96 | 5.44 | 5.94 | 6.46 | 6.83 | 7.04 | 7.16 | 7.23 |
| 130 | 4.58 | 4.76 | 5.12 | 5.60 | 6.11 | 6.59 | 6.92 | 7.10 | 7.20 | 7.26 |
| 140 | 4.72 | 4.90 | 5.26 | 5.75 | 6.24 | 6.68 | 6.98 | 7.15 | 7.23 | 7.27 |
| 150 | 4.87 | 5.06 | 5.44 | 5.87 | 6.36 | 6.78 | 7.04 | 7.18 | 7.25 | 7.29 |

VOLUME PER ACRE IN BOARD FEET

Volumes in board feet of all trees 8 inches and larger in diameter, for stands of average stocking and composition, are given in Table 13 and Figure 8. The log rule used is the International, 1/8-inch kerf, and the volume is of that part of the stem between a stump 1 foot high and a top diameter inside bark of 5 inches.

TABLE 13.—Volume per acre in board feet (International rule, 1/8-inch kerf), average stands

| Age (years) | Board feet per acre on site index— | | | | | | | | | |
|-------------|------------------------------------|--------|--------|--------|--------|---------|---------|---------|---------|---------|
| | 25 | 30 | 40 | 50 | 60 | 70 | 80 | 90 | 100 | 110 |
| 30..... | | | 85 | 450 | 1,100 | 2,120 | 3,840 | 6,030 | 8,670 | 11,110 |
| 40..... | 700 | 940 | 1,740 | 3,210 | 5,540 | 9,100 | 14,300 | 20,500 | 26,800 | 33,450 |
| 50..... | 2,300 | 2,950 | 5,050 | 8,300 | 13,200 | 20,340 | 29,850 | 40,100 | 50,500 | 60,700 |
| 60..... | 4,720 | 6,100 | 9,970 | 15,750 | 23,500 | 34,300 | 47,900 | 61,950 | 76,750 | 90,200 |
| 70..... | 8,330 | 10,250 | 15,950 | 24,300 | 35,400 | 48,950 | 65,400 | 83,250 | 102,100 | 118,900 |
| 80..... | 11,900 | 14,550 | 21,950 | 32,700 | 45,300 | 61,650 | 81,400 | 102,400 | 122,800 | 142,300 |
| 90..... | 15,450 | 18,500 | 27,950 | 40,000 | 54,300 | 73,000 | 95,750 | 118,400 | 142,200 | 161,900 |
| 100..... | 18,600 | 22,300 | 33,100 | 46,850 | 63,150 | 84,250 | 108,200 | 133,400 | 158,200 | 179,400 |
| 110..... | 21,550 | 25,900 | 38,100 | 53,450 | 70,700 | 93,850 | 120,100 | 146,200 | 172,200 | 195,600 |
| 120..... | 24,600 | 29,250 | 42,400 | 58,750 | 78,400 | 103,400 | 130,500 | 158,400 | 186,200 | 209,700 |
| 130..... | 27,000 | 31,900 | 46,350 | 64,400 | 85,550 | 111,400 | 140,500 | 170,400 | 198,000 | 222,900 |
| 140..... | 29,350 | 34,650 | 50,250 | 69,600 | 92,050 | 118,600 | 149,400 | 180,200 | 209,700 | 235,500 |
| 150..... | 31,900 | 37,600 | 54,400 | 74,550 | 98,250 | 125,800 | 157,700 | 189,600 | 220,400 | 247,600 |

This table was constructed in the usual manner, by multiplying cubic-foot volume for any age and site index by the board foot-cubic foot ratio corresponding to the average diameter for that age and site index.

This average table is corrected for specific stocking and composition in a similar manner. The corrected average diameter is first determined, and the corresponding board foot-cubic foot ratio is then multiplied by the corrected volume in cubic feet, giving the corrected volume in board feet.

MEAN ANNUAL INCREMENT

Table 14 and Figure 9 show the mean annual increment in cubic feet, for stands of average stocking and composition. These values were derived by dividing the values in Table 10 by the corresponding ages. Mean annual increment culminates at 70 years for any site quality.

TABLE 14.—Mean annual growth in cubic feet per acre, average stands

| Age (years) | Mean annual growth in cubic feet per acre on site index— | | | | | | | | | |
|-------------|--|-------|-------|-------|--------|--------|--------|--------|--------|--------|
| | 25 | 30 | 40 | 50 | 60 | 70 | 80 | 90 | 100 | 110 |
| 30..... | 31.00 | 35.33 | 47.33 | 60.00 | 73.33 | 88.33 | 106.67 | 125.67 | 146.00 | 161.67 |
| 40..... | 39.75 | 45.00 | 60.25 | 77.25 | 94.25 | 113.75 | 137.50 | 162.50 | 186.25 | 211.25 |
| 50..... | 45.20 | 51.80 | 69.60 | 88.80 | 108.40 | 131.20 | 158.80 | 186.00 | 214.00 | 240.00 |
| 60..... | 49.17 | 55.83 | 75.17 | 96.50 | 117.00 | 142.50 | 171.67 | 201.67 | 231.67 | 258.33 |
| 70..... | 50.00 | 57.14 | 76.43 | 97.86 | 120.71 | 145.71 | 174.29 | 204.29 | 235.71 | 262.86 |
| 80..... | 50.00 | 56.75 | 76.25 | 97.50 | 119.00 | 143.75 | 172.50 | 202.50 | 232.50 | 261.25 |
| 90..... | 49.11 | 55.56 | 75.00 | 95.56 | 116.67 | 141.11 | 168.89 | 197.78 | 230.00 | 256.67 |
| 100..... | 48.30 | 55.00 | 74.00 | 94.50 | 115.00 | 139.00 | 166.00 | 195.00 | 226.00 | 252.00 |
| 110..... | 47.27 | 54.09 | 72.73 | 92.73 | 111.82 | 135.45 | 162.73 | 190.91 | 220.45 | 247.27 |
| 120..... | 46.67 | 53.25 | 71.25 | 90.00 | 110.00 | 133.33 | 159.17 | 187.50 | 216.67 | 241.67 |
| 130..... | 45.38 | 51.54 | 69.62 | 88.46 | 107.69 | 130.00 | 156.15 | 184.62 | 211.54 | 236.15 |
| 140..... | 44.43 | 50.50 | 68.21 | 86.43 | 105.36 | 126.79 | 152.86 | 180.00 | 207.14 | 231.43 |
| 150..... | 43.67 | 49.53 | 66.67 | 84.67 | 103.00 | 123.67 | 149.33 | 176.00 | 202.67 | 226.67 |




FIGURE 9.—Mean annual increment in cubic-foot volume culminates at 70 years on all sites

Table 15 and Figure 10 show the mean annual increment in board feet. The age of culmination varies with site quality, from 95 years for site index 110 to over 150 years for site index 30. For site index 70 the age of culmination is about 120 years.

TABLE 15.—Mean annual growth in board feet per acre (international rule, 1/8-inch kerf), average stands

| Age (years) | Mean annual growth in board feet per acre on site index— | | | | | | | | | |
|-------------|--|-------|-------|-------|-------|-------|---------|---------|---------|---------|
| | 25 | 30 | 40 | 50 | 60 | 70 | 80 | 90 | 100 | 110 |
| 30..... | | | 2.8 | 15.0 | 36.7 | 70.7 | 128.0 | 201.0 | 289.0 | 370.3 |
| 40..... | 17.5 | 23.5 | 43.5 | 80.2 | 135.5 | 227.5 | 357.5 | 512.5 | 670.0 | 836.2 |
| 50..... | 46.0 | 59.6 | 101.0 | 166.0 | 264.0 | 406.8 | 597.0 | 802.0 | 1,010.0 | 1,214.0 |
| 60..... | 78.7 | 101.7 | 166.2 | 262.5 | 391.7 | 571.7 | 798.3 | 1,032.5 | 1,279.2 | 1,503.3 |
| 70..... | 119.0 | 146.4 | 227.9 | 347.1 | 505.7 | 699.3 | 934.3 | 1,189.3 | 1,458.6 | 1,698.6 |
| 80..... | 148.8 | 181.9 | 274.4 | 408.8 | 566.2 | 770.6 | 1,017.5 | 1,280.0 | 1,535.0 | 1,768.8 |
| 90..... | 171.7 | 205.6 | 310.6 | 444.4 | 603.3 | 811.1 | 1,063.9 | 1,315.6 | 1,580.0 | 1,778.5 |
| 100..... | 186.0 | 223.0 | 331.0 | 468.5 | 631.5 | 842.5 | 1,082.0 | 1,334.0 | 1,582.0 | 1,794.0 |
| 110..... | 195.9 | 235.5 | 346.4 | 485.9 | 642.7 | 853.2 | 1,091.3 | 1,329.1 | 1,565.5 | 1,778.2 |
| 120..... | 205.0 | 243.8 | 353.3 | 489.6 | 653.3 | 861.7 | 1,087.5 | 1,320.0 | 1,551.7 | 1,747.5 |
| 130..... | 207.7 | 245.4 | 356.5 | 495.4 | 658.1 | 856.9 | 1,080.8 | 1,310.8 | 1,523.1 | 1,714.6 |
| 140..... | 209.6 | 247.5 | 358.9 | 497.1 | 657.5 | 847.1 | 1,067.1 | 1,287.1 | 1,497.9 | 1,682.1 |
| 150..... | 212.7 | 250.7 | 362.7 | 497.0 | 655.0 | 838.7 | 1,051.3 | 1,264.0 | 1,469.3 | 1,652.7 |




FIGURE 10.—Mean annual increment in board feet culminates at 95 years on the best sites and at 150 years and more on the poorest sites

SUMMARY

With the continued depletion of old-growth stands in the very important mixed conifer type of California there is increasing need for adequate tables of the second-growth timber that is rapidly coming in on many cut-over areas in this region. The tables here presented are based in the main on intensive studies of 311 sample plots which were selected as supplying reasonably representative, although inevitably incomplete data. The position, range, and occurrence of the second-growth mixed conifer stands is described as well as the conditions for their establishment and the relative importance of virgin and second-growth forest in this type.

In view of the number of unusual features that were involved in this yield study a careful explanation of the technic employed in the preparation of the tables is included as an appendix.

LITERATURE CITED

- (1) BAKER, F. S.
1923. NOTES ON THE COMPOSITION OF EVEN-AGED STANDS. Jour. Forestry 21: 712-717, illus.
- (2) BRUCE, D.
1923. PRELIMINARY YIELD TABLES FOR SECOND-GROWTH REDWOOD. Calif. Agr. Expt. Sta. Bul. 361, p. 425-467, illus.
- (3) ———
1926. A METHOD OF PREPARING TIMBER-YIELD TABLES. Jour. Agr. Research 32: 543-557, illus.
- (4) BRUCE D. and REINEKE, L. H.
1929. THE USE OF ALINEMENT CHARTS IN CONSTRUCTING FOREST STAND TABLES. Jour. Agr. Research 38: 289-308, illus.
- (5) HAIG, I. T.
1932. SECOND-GROWTH YIELD, STAND, AND VOLUME TABLES FOR THE WESTERN WHITE PINE TYPE. U. D. Dept. Agr. Tech. Bul. 323, 68 p., illus.
- (6) MEYER, W. H.
1930. DIAMETER DISTRIBUTION SERIES IN EVEN-AGED FOREST STANDS. Yale Univ. School Forestry Bul. 28, 105 p., illus.
- (7) REINEKE, L. H.
1927. A MODIFICATION OF BRUCE'S METHOD OF PREPARING TIMBER-YIELD TABLES. Jour. Agr. Research 35: 843-856, illus.
- (8) SCHUMACHER, F. X.
1926. YIELD, STAND, AND VOLUME TABLES FOR WHITE FIR IN THE CALIFORNIA PINE REGION. Calif. Agr. Expt. Sta. Bul. 407, 26 p., illus.

APPENDIX

SPECIAL TECHNIC USED IN THE PREPARATION OF THE TABLES

SITE INDEX

The mixed-conifer types present a problem in the determination of site quality by reason of their variation in composition. If dominant trees of a given species were invariably present, the height of trees of that species alone could be used as the index of site quality. Since this is not the case, the relative height growths of the various species must be compared, to determine which species it is desirable to use and to what extent the heights of each depart from the heights of the other species.

The actual analysis showed a simple solution for the problem. The height of the average dominant tree of each species on a plot was first determined. Using all plots for which such values could be determined for two or more species, it was found that white fir was common to nearly all. Accordingly, dominant heights of other species on each plot were expressed as a percentage of the dominant height of white fir. The percentages so derived were then plotted over age, by species. The resultant curves were horizontal straight lines and passed through 98 per cent for ponderosa pine and Douglas fir, 100 per cent for red fir, and 90 per cent for sugar pine. (No data were available for incense cedar, rarely dominant.) This is in close agreement with the values obtained by Schumacher (8) and by Bruce (2).

These results indicate that site index can be determined directly from white fir, ponderosa pine, Douglas fir, and red fir, alone or in any combination. Naturally, the use of as many of these species as are present will give a more accurate value of site index.

STAND-DENSITY INDEX

In any even-aged stand, the curve of number of trees by breast-high diameter classes has a definite characteristic form, often approaching that of the normal frequency curve (1, 4, 6). Such a curve can be mathematically described in several ways, a description commonly used being the statement of the mean or average diameter and the standard deviation (3). For even-aged stands, however, the numerical value of the standard deviation is positively correlated with the average diameter. Therefore, average diameter alone may be used to describe stands of similar frequency-curve form where extreme accuracy is not needed.

If, then, a large number of stands of the same description, that is, of the same average diameter, be compared as to number of trees per acre, it is obvious that the stand with most trees is the most fully stocked. If this be taken to represent

the maximum or complete stocking, the number of trees in each of these stands expressed as a percentage of this maximum will give the percentage stocking of the stand.

The number of trees per acre for complete stocking varies with the average diameter of the stand. The curve representing the relationship between number of trees per acre and average diameter corresponds (8) to the formula

$$\log \text{ number of trees} = -1.605 \log \text{ average diameter breast high} + k$$

where k is a constant varying with species. This formula plots as a straight line on logarithmic graph paper (the solid line of Figure 2).

Since the constant k differs with species, it would be impractical to determine its value for the innumerable mixtures of the mixed-conifer types, nor would differences for the common proportions of those species be sufficiently great to justify the attempt. For simplicity then, k was taken as 4.605 to give a curve passing through 1,000 trees per acre at 10 inches average diameter. This curve is used as a reference curve; its elevation is of no importance aside from the convenience of its round-number coordinates (10, 1,000).

The reference curve, thus arbitrarily chosen, is somewhat high for the mixed-conifer stands of this study, though it is not too high for stands of certain other species. It was deemed more desirable, however, to refer stocking to a standard curve suitable for many species than to a curve applicable only to the species of this study. Since this reference curve is higher than the curve representing maximum stocking for the mixed-conifer types, percentage stocking values based on it will be higher than the true percentage stocking. Although this distortion could be easily cared for in any computations involved, it would be somewhat confusing in making mental comparisons. For this reason, the stand-density index was devised.

Any curve representing a constant percentage of the reference curve will plot as a straight line parallel to the reference curve. The curves corresponding to all possible percentages thus constitute an infinite series of parallel curves. Since their slopes are identical, the position of any curve of the series may be defined by any one ordinate and abscissa. If the same abscissa be used for all curves, each curve may be defined by ordinate alone. Accordingly, an abscissa of 10 inches average diameter was chosen, and the ordinate (number of trees per acre) corresponding to this abscissa is termed "stand-density index."

Stand-density index, as here conceived, is obviously independent of species variation in elevation of the curve of maximum number of trees, since it is an absolute, not a relative measure.

The effect of age and site quality on the relationship between number of trees and average diameter is very small. Multiple correlation coefficients for various species ranged from 0.141 ± 0.050 to 0.251 ± 0.049 . These are not significant. The corresponding alienation coefficients were 0.990 and 0.968, respectively. For all practical purposes, therefore, no correction need be made for age or site quality.

ADJUSTMENT OF PLOT VALUES FOR STOCKING IN CONSTRUCTION OF TABLES FOR AVERAGE STOCKING

Having a measure of stocking, it should be possible to eliminate the effects of variation in stocking by converting individual plot values to a uniform-stocking basis, thus reducing the dispersion of the plotted points and eliminating the effect of any irregular distribution of stocking. As a result the required curves of the various stand measures for average stocking and composition should be well defined and easily fitted.

The individual plot values for total basal area, number of trees, and cubic-foot volume were accordingly modified by straight proportion between plot stand-density index and average stand-density index (479). Thus, the values for a plot of stand-density index 700 were reduced by the proportion of 700 to 479, and a plot of stand-density index 450 increased by the proportion 450 to 479. The usual pairs of average curves (7) such as total basal area over age and percentage of average total basal area over site index, were fitted to the modified values. As expected, the dispersion of the values thus derived was reduced materially, and the trends of the curves were more consistent and better defined.

Since modification for stocking was effected by increasing or decreasing the number of trees of a given average diameter, the average diameter values of the individual plots were not changed. The average curves were based on these original plot values of diameter and checked through the modified total basal area and modified number of tree curves; the agreement between them was very close.

The board-foot tables are derived from the cubic-foot tables and the curve of board foot-cubic foot ratio; board-foot values, therefore, were not modified for stocking.

ANALYSIS OF EFFECT OF COMPOSITION AND STOCKING

By using the table of total basal area for stands of average stocking and composition, the tabular value corresponding to the age and site index of each sample plot was determined. The basal area of each plot was then expressed as a percentage of this tabular value. These percentages were used as the dependent-variable values in a multiple linear correlation, with percentage stocking and composition by species as the independent variables. The resulting multiple-regression equation was—

Total basal area per cent
 = 2.0303 (stand-density index) + 0.1493 (ponderosa pine per cent)
 + 0.1545 (sugar pine per cent) + 0.0541 (Douglas fir per cent) + 0.0980
 (white fir per cent) + 0.1229 (incense cedar per cent) + 0.1883 (red
 fir per cent) - 8.1511.

The regression coefficients, in combination with the numerical values of the variables, indicate that variations in stocking account for the greatest part of the variation in basal area, the effect of composition being relatively small.

A similar correlation was made for number of trees per acre, resulting in the regression equation—

Number of trees per cent
 = 2.0058 (stand-density index) - 0.3166 (ponderosa pine per cent)
 - 0.2973 (sugar pine per cent) - 0.0159 (Douglas fir per cent) - 0.1355
 (white fir per cent) - 0.1358 (incense cedar per cent) - 0.0905 (red
 fir per cent) + 19.3572.

• Here, also, the greatest effect is assignable to variation in stand density.

A third correlation was made, in a similar manner, for volume in cubic feet. The resulting regression equation was—

Cubic-foot volume per cent
 = 1.5251 (stand-density index) - 0.7170 (ponderosa pine per cent)
 - 0.7740 (sugar pine per cent) - 0.9398 (Douglas fir per cent) - 1.0704
 (white fir per cent) - 0.8427 (incense cedar per cent) - 1.1996 (red
 fir per cent) + 128.7553.

STATISTICAL MEASURES

The various statistical measures employed in the computations are given in Table 16.

TABLE 16.—Statistical measures

| Dependent variable (per cent) | Aliena- tion index | Correla- tion index | Standard error of average table | Standard error of average table and regres- sion equation combined | Average percentage deviation ¹ | |
|-------------------------------|--------------------------|---------------------------|--|---|--|---|
| | | | | | From average table only | From average table and regres- sion equation combined |
| Total basal area..... | 0.2486 | 0.9686 | <i>Per cent</i> 26.24 | <i>Per cent</i> 6.52 | <i>Per cent</i> 21.0 | <i>Per cent</i> 5.22 |
| Number of trees..... | .6926 | .7213 | 32.19 | 22.29 | 25.8 | 17.8 |
| Volume, cubic feet..... | .7442 | .6680 | 27.52 | 20.48 | 22.0 | 16.4 |
| Volume, board feet..... | .7224 | .6915 | 36.46 | 26.34 | 29.2 | 21.1 |

¹ Average percentage deviation = $\frac{\text{standard error}}{1.25}$